


Deprescribing guideline implementation experiences

Barbara Farrell BScPhm, PharmD, FCSHP
Scientist, Bruyère Research Institute
Ottawa, Canada
March 27, 2018







Developing and implementing deprescribing guidelines for the elderly HRSF grant (2013-2016)



Findings

- There is an appetite for such guidelines; 14 priorities identified
- Deprescribing decision-support algorithms easily implemented into routine pharmacist-physician LTC medication reviews; appeared to increase self-efficacy for deprescribing and reduce target medication use
- In LTC, need better patient/family/staff buy-in (and communication)
- Implementation in Family Health Teams more challenging due to competing priorities, EMR limitations + lack of documented reason



Catalyst grant (2016-2017)



Canadian Foundation
for Pharmacy

- Objectives
 - Explore and build community pharmacists' capacity to integrate deprescribing into workflow
 - Quantify deprescribing opportunities, actions and outputs
 - Develop a viable pharmacy business model to integrate deprescribing into pharmacy practice
- Methods
 - Four community pharmacies + Advisory Group
 - Education and resources provided
 - Quantify opportunities for deprescribing, describe activities and processes associated with guideline use; including how long such activities take, and highlight enablers and barriers encountered
 - Use iterative observation periods (PDSA cycles) in each pharmacy and discussion of findings with the Advisory Group to hone workflow strategies over time, and to provide motivation and build capacity for scale up of the practice of deprescribing



deprescribing.org

INSTITUT DE RECHERCHE

Bruyère
RESEARCH INSTITUTE

Findings

- Deprescribing was feasible at all sites
 - All sites able to identify individual and common goals, and develop unique community pharmacy workflow models for deprescribing
- Deprescribing practices differed by site
- Each pharmacy developed resources and materials at their sites to integrate deprescribing
- 4 common deprescribing steps noted across sites
 - Capacity building activities
 - Preliminary interactions
 - Detailed interactions with the pharmacist
 - Follow-up and monitoring



Findings

Facilitators

- Supportive staff and students that were motivated regarding deprescribing
- On-site educational initiatives
- Approaches to draw people into the pharmacy
- Employing a collaborative team approach (all staff trained)
- Enhancing patients' awareness and education regarding the risks and options to reassess
- The development of standard templates to reduce time spent on each Pharmaceutical Opinion
- Faxing algorithms along with PO

Challenges

- Competing workload and time
- Staff turnover and new staff training
- Communication delays and lack of response from prescribers
- Patients uncertain about change
- Patients receiving medications by delivery and/or using multiple pharmacies
- Duplication of documentation
- Inadequate compensation models for time required
- Workspace limitations



Next

- Community engagement for deprescribing initiatives (DICE) – HSRF 2017-2020



deprescribing.org

INSTITUT DE RECHERCHE
Bruyère
RESEARCH INSTITUTE

Evidence Based Deprescribing Guideline Symposium 2018

Supported By:



CENTRE FOR AGING
+ BRAIN HEALTH
INNOVATION
Powered by Baycrest



Sponsored By:



Canadian
Deprescribing
Network



MEDICAL
PHARMACIES

CADTH

Evidence
Driven.

Canadian Foundation for **Healthcare Improvement**

Fondation canadienne pour **l'amélioration des services de santé**