

Why is patient taking a BZRA?

If unsure, find out if history of anxiety, past psychiatrist consult, whether may have been started in hospital for sleep, or for grief reaction.

- Insomnia on its own OR insomnia where underlying comorbidities managed
For those 65 years of age: taking BZRA regardless of duration (avoid as first line therapy in older people)
For those 18-64 years of age: taking BZRA > 4 weeks

- Other sleeping disorders (e.g. restless legs)
- Unmanaged anxiety, depression, physical or mental condition that may be causing or aggravating insomnia
- Benzodiazepine effective specifically for anxiety
- Alcohol withdrawal

Engage patients (discuss potential risks, benefits, withdrawal plan, symptoms and duration)

Recommend Deprescribing

Continue BZRA

Taper and then stop BZRA

(taper slowly in collaboration with patient, for example ~25% every two weeks, and if possible, 12.5% reductions near end and/or planned drug-free days)

- For those ≥ 65 years of age (strong recommendation from systematic review and GRADE approach)
- For those 18-64 years of age (weak recommendation from systematic review and GRADE approach)
- Offer behavioural sleeping advice; consider CBT if available (see reverse)

- Minimize use of drugs that worsen insomnia (e.g. caffeine, alcohol etc.)
- Treat underlying condition
- Consider consulting psychologist or psychiatrist or sleep specialist

Monitor every 1-2 weeks for duration of tapering

Expected benefits:

- May improve alertness, cognition, daytime sedation and reduce falls

Withdrawal symptoms:

- Insomnia, anxiety, irritability, sweating, gastrointestinal symptoms (all usually mild and last for days to a few weeks)

Use non-drug approaches to manage insomnia
Use behavioral approaches and/or CBT (see reverse)

If symptoms relapse:
Consider

- Maintaining current BZRA dose for 1-2 weeks, then continue to taper at slow rate

Alternate drugs

- Other medications have been used to manage insomnia. Assessment of their safety and effectiveness is beyond the cope of this algorithm. See BZRA deprescribing guideline for details.

© Use freely, with credit to the authors. Not for commercial use. Do not modify or translate without permission.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. Contact deprescribing@bruyere.org or visit deprescribing.org for more information.

This algorithm and accompanying advice support recommendations in the NICE guidance on the use of zaleplon, zolpidem and zopiclone for the short-term management of insomnia, and medicines optimisation. **National Institute for Health and Care Excellence, February 2019**



BZRA Availability

| BZRA | Strength |
|---|-----------------------------|
| Alprazolam (Xanax®) ^T | 0.25 mg, 0.5 mg, 1 mg, 2 mg |
| Bromazepam (Lectopam®) ^T | 1.5 mg, 3 mg, 6 mg |
| Chlordiazepoxide ^C | 5 mg, 10 mg, 25 mg |
| Clonazepam (Rivotril®) ^T | 0.25 mg, 0.5 mg, 1 mg, 2 mg |
| Clorazepate (Tranxene®) ^C | 3.75 mg, 7.5 mg, 15 mg |
| Diazepam (Valium®) ^T | 2 mg, 5 mg, 10 mg |
| Flurazepam (Dalmane®) ^C | 15 mg, 30 mg |
| Lorazepam (Ativan®) ^{T,S} | 0.5 mg, 1 mg, 2 mg |
| Nitrazepam (Mogadon®) ^T | 5 mg, 10 mg |
| Oxazepam (Serax®) ^T | 10 mg, 15 mg, 30 mg |
| Temazepam (Restoril®) ^C | 15 mg, 30 mg |
| Triazolam (Halcion®) ^T | 0.125 mg, 0.25 mg |
| Zopiclone (Imovane®, Rhovane®) ^T | 5 mg, 7.5 mg |
| Zolpidem (Sublinox®) ^S | 5 mg, 10 mg |

T = tablet , C = capsule, S = sublingual tablet

BZRA Side Effects

- BZRAs have been associated with:
 - physical dependence, falls, memory disorder, dementia, functional impairment, daytime sedation and motor vehicle accidents
- Risks increase in older persons

Engaging patients and caregivers

Patients should understand:

- The rationale for deprescribing (associated risks of continued BZRA use, reduced long-term efficacy)
- Withdrawal symptoms (insomnia, anxiety) may occur but are usually mild, transient and short-term (days to a few weeks)
- They are part of the tapering plan, and can control tapering rate and duration

Tapering doses

- No published evidence exists to suggest switching to long-acting BZRAs reduces incidence of withdrawal symptoms or is more effective than tapering shorter-acting BZRAs
- If dosage forms do not allow 25% reduction, consider 50% reduction initially using drug-free days during latter part of tapering, or switch to lorazepam or oxazepam for final taper steps

Behavioural Management

Primary care:

- Go to bed only when sleepy
- Do not use bed or bedroom for anything but sleep (or intimacy)
- If not asleep within about 20-30 min at the beginning of the night or after an awakening, exit the bedroom
- If not asleep within 20-30 min on returning to bed, repeat #3
- Use alarm to awaken at the same time every morning
- Do not nap
- Avoid caffeine after noon
- Avoid exercise, nicotine, alcohol, and big meals within 2 hrs of bedtime

Institutional care:

- Pull up curtains during the day to obtain bright light exposure
- Keep alarm noises to a minimum
- Increase daytime activity & discourage daytime sleeping
- Reduce number of naps (no more than 30 mins and no naps after 2 pm)
- Offer warm decaf drink, warm milk at night
- Restrict food, caffeine, smoking before bedtime
- Have the resident toilet before going to bed
- Encourage regular bedtime and rising times
- Avoid waking at night to provide direct care
- Offer backrub, gentle massage

Using CBT

What is cognitive behavioural therapy (CBT)?

- CBT includes 5-6 educational sessions about sleep/insomnia, stimulus control, sleep restriction, sleep hygiene, relaxation training and support

Does it work?

- CBT has been shown in trials to improve sleep outcomes with sustained long-term benefits

Who can provide it?

- Clinical psychologists usually deliver CBT, however, others can be trained or can provide aspects of CBT education; self-help programs are available

How can providers and patients find out about it?

- Some resources can be found here: <http://sleepwellns.ca/>

© Use freely, with credit to the authors. Not for commercial use. Do not modify or translate without permission.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. Contact deprescribing@bruyere.org or visit deprescribing.org for more information.

Pottie K, Thompson W, Davies S, Grenier J, Sadowski C, Welch V, Holbrook A, Boyd C, Swenson JR, Ma A, Farrell B. Evidence-based clinical practice guideline for deprescribing benzodiazepine receptor agonists. Can Fam Physician 2018;64:339-51 (Eng), e209-24 (Fr)

This algorithm and accompanying advice support recommendations in the NICE guidance on the use of zaleplon, zolpidem and zopiclone for the short-term management of insomnia, and medicines optimisation. National Institute for Health and Care Excellence, February 2019



deprescribing.org

